



PROVIRAPLEX



[HOME](#) | [SPECIALS](#) | [NEW PRODUCTS](#) | [ALL PRODUCTS](#) | [REVIEWS](#) | [CONTACT US](#) | [FAQ](#)

PROVIRAPLEX

PROVIRAPLEX (Mesterolone) Axiolabs BUY PROVIRAPLEX

Categories

- [Injectable Steroids](#)
- [Oral Steroids](#)
- [Ancillaries](#)
- [Sexual Health](#)
- [Syringes & Needles](#)

News

2008-06-17

[Anabolic Steroid Dianabol Turns 50](#)

The performance-enhancing drug Dianabol, which became the first widely used anabolic steroid and has built muscle for athletes and bodybuilders and become the subject of criminal prosecutions, was developed in 1958. There's a good piece in the New York Daily News looking at half a century of the steroid's use:

2008-06-27

[Women On Prudent Dietary Pattern May Reduce Risk Of Death](#)


Women On Prudent Dietary Pattern May Reduce Risk Of Death Women who eat a diet rich in vegetables, fruits, legumes, whole grains, fish and poultry may reduce their risk of death from cardiovascular disease and all causes. Women who follow a traditional "Western" diet of red and processed meat, refined grains, fries and sweets may increase their risk. That's the

<http://proviraplex.net/> (1 of 2)6/29/2008 8:38:19 AM

Substance: Mesterolone
Manufactured by: Axiolabs
Packaging: 20 tabs, each tab 25mg

Proviraplex (Mesterolone) is an androgenic/anabolic steroid with a particularly high affinity for SHBG (sex hormone binding globulin). SHBG is a protein that acts by binding to sex hormones and preventing them from interacting within the body at places like the AR (androgen receptor). Because of its high affinity with SHBG, Proviraplex' main use is to competitively bind with SHBG, freeing up more of the endogenous hormone to be active within the body. In effect, proviron acts as a 'magnifier' of the anabolic effects of steroid hormones. It also exhibits anti-aromatase activity, hence helping to prevent the conversion of certain steroids into estrogen. As a result, the intake of proviraplex whilst on a cycle of steroids, results in a 'hardening' of skeletal muscle - a look which is very popular amongst steroid users.

One side effect of proviraplex is the increased frequency and severity of erections, particularly during sleep/upon waking. Although proviraplex is an orally administered steroid, and is alkylated to survive the first pass through the liver, it is considered to be almost completely non-toxic to the liver. A typical dose of proviron might be from 25-100mg daily.



Proviraplex (Proviron) to Tamoxifen.

With **Proviraplex (Proviron)** the athlete obtains more muscle hardness since the androgen level is increased and the estrogen concentration remains low. This, in particular, is noted positively during the preparation for a competition when used in combination with a diet. Female athletes who naturally have a higher estrogen level often supplement their steroid intake with **Proviron** resulting in an increased muscle hardness. In the past it was common for bodybuilders to take a daily dose of one 25 mg tablet over several weeks, sometimes even months, in order to appear hard all year round.

This was especially important for athletes appearances at guest performances, seminars and photo sessions. Today Clenbuterol is usually taken over the entire year since possible virilization symptoms cannot occur which is not yet the case with Proviron. Since Proviron is very effective male athletes usually need only 50 mg/day which means that the athlete usually takes one 25 mg tablet in the morning and another 25 mg tablet in the evening. In some cases one 25 mg tablet per day is sufficient. When combining **Proviraplex (Proviron)** with Tamoxifen (50 mg Proviron/day and 20 mg Tamoxifen/day) this will lead to an almost complete suppression of estrogen.

The side effects of **Proviraplex (Proviron)** in men are low at a dosage of 2-3 tablets/day so that Proviron, taken for example in combination with a steroid cycle, can be used comparatively without risk over several weeks. Since Proviron is well-tolerated by the liver liver dysfunctions do not occur in the given dosages. For athletes who are used to acting under the motto "more is better" the intake of Proviron could have a paradoxical effect.

[Buy Now](#)



conclusion of researchers who reported the results of a Harvard School of Public Health study in Circulation: Journal of the American Heart Association. The study of 72,113 healthy women found that high adherence to the "prudent" dietary pattern was associated with a 28 percent lower risk of death from cardiovascular disease and a 17 percent lower risk of premature death from all causes when compared to the lowest adherence.

The most common side effect of Proviron-or in this case, secondary symptom- is in part a distinct sexual overstimulation and in some cases continuous penis erection. Since this condition can be painful and lead to possible damages, a lower dosage or discontinuing the compound are the only sensible solutions. Female athletes should use Proviron® with caution since possible androgenic side effects cannot be excluded. Women who want to give Proviron a try should not take more than one 25 mg tablet per day. Higher dosages and periods of intake of more than four weeks considerably increase the risk of virilization symptoms. Female athletes who have no difficulties with Proviron® obtain good results with 25 mg Proviron/day and 20 mg Tamoxifen/day and, in combination with a diet, report an accelerated fat breakdown and continuously harder muscles.

[Home](#)|[Specials](#)|[New Products](#)|[All Products ...](#)|[Reviews](#)|[Contact Us](#)|[FAQ](#)|[Map](#)

BUY PROVIRAPLEX Copyright © 2005-2007. [Privacy Notice](#)