



SILDENAPLEX 100



[HOME](#) | [SPECIALS](#) | [NEW PRODUCTS](#) | [ALL PRODUCTS](#) | [REVIEWS](#) | [CONTACT US](#) | [FAQ](#)

SILDENAPLEX 100

SILDENAPLEX 100 (Sildenafil citrate) Axiolabs

Categories

- [Injectable Steroids](#)
- [Oral Steroids](#)
- [Ancillaries](#)
- [Sexual Health](#)
- [Syringes & Needles](#)

News

2008-06-27

[Women On Prudent Dietary Pattern May Reduce Risk Of Death](#)

Women On Prudent Dietary Pattern May Reduce Risk Of Death Women who eat a diet rich in vegetables, fruits, legumes, whole grains, fish and poultry may reduce their risk of death from cardiovascular disease and all causes. Women who follow a traditional "Western" diet of red and processed meat, refined grains, fries and sweets may increase their risk. That's the conclusion of researchers who reported the results of a Harvard School of Public Health study in Circulation: Journal of the American Heart Association. The study of 72,113 healthy women found that high adherence to the "prudent" dietary pattern was associated with a 28 percent lower risk of death from cardiovascular disease and a 17 percent lower risk of premature death from all causes when compared to the lowest adherence.

Substance: Sildenafil Citrate
Manufactured by: Axiolabs
Packaging: 4 tabs, each tab 100mg

Sildenafil 100 (VIAGRA) is an oral medication for erectile dysfunction (ED). It helps most men with ED improve their erections. That means they are able to achieve harder erections.

Getting an erection is only half the story. Maintaining an erection is just as important as getting one. **VIAGRA** can help with both. This can lead to a more satisfying sexual experience with your partner.

Who can take **VIAGRA**


VIAGRA has an established safety profile of over 9 years. It has been shown to work safely and effectively in men with ED. This includes:

- [Men of all ages](#)
- [Men who have difficulty all the time or just some of the time](#)
- [Men with other health issues such as high blood pressure, diabetes, and depression](#)

Find out if **VIAGRA** is right for you. Speak to your doctor. He can tell you if you have ED. And can recommend an effective treatment.

DOSAGE AND ADMINISTRATION

For most patients, the recommended dose is 50 mg taken, as needed, approximately 1 hour before sexual activity. However, **VIAGRA** may be taken anywhere from 4 hours to 0.5 hour before sexual activity. Based on effectiveness and toleration, the dose may be increased to a maximum recommended dose of 100 mg or decreased to 25 mg. The maximum recommended dosing frequency is once per day.



VIAGRA in a 48 hour period.

VIAGRA was shown to potentiate the hypotensive effects of nitrates and its administration in patients who use nitric oxide donors or nitrates in any form is therefore contraindicated. When **VIAGRA** is co-administered with an alpha-blocker, patients should be stable on alphablocker

[Buy Now](#)



The following factors are associated with increased plasma levels of sildenafil: age >65 (40% increase in AUC), hepatic impairment (e.g., cirrhosis, 80%), severe renal impairment (creatinine clearance <30 mL/min, 100%), and concomitant use of potent cytochrome P450 3A4 inhibitors [ketoconazole, itraconazole, erythromycin (182%), saquinavir (210%)]. Since higher plasma levels may increase both the efficacy and incidence of adverse events, a starting dose of 25 mg should be considered in these patients.

Ritonavir greatly increased the systemic level of sildenafil in a study of healthy, non-HIV infected volunteers (11-fold increase in AUC, see Drug Interactions.) Based on these pharmacokinetic data, it is recommended not to exceed a maximum single dose of 25 mg of

2008-06-27

[The sport of bodybuilding](#)

The sport of bodybuilding The popular, growing sport of bodybuilding in the region is leading to bigger championships for local enthusiasts. Dana Ali, 31, who recently returned to Kurdistan from Britain, started lifting weights six year ago when he was abroad. "Oh, I became many pounds overweight," he said, taking a break after finishing the first set. "I see so many [overweight people] here, and I don't want to be like them. "I feel better and forget about all my problems when I come in and exercise," said Ali, having arrived straight from work wearing a black T-shirt and leather gloves. He works out five times a week and has shed most of his body fat; his chest and biceps are as thick as they were when he was living in Britain.

therapy prior to initiating **VIAGRA** treatment and **VIAGRA** should be initiated at the lowest dose (see Drug Interactions).

Side effects

Amongst sildenafil's rare but serious adverse effects are: priapism, severe hypotension, myocardial infarction, ventricular arrhythmias, stroke and increased intraocular pressure.

Common side effects include sneezing, headache, flushing, dyspepsia, prolonged erections, palpitations and photophobia. Visual changes including blurring of vision and a curious bluish tinge have also been reported.

Care should be exercised by patients who are also taking Protease inhibitors for the treatment of HIV. Protease inhibitors inhibit the metabolism of sildenafil, effectively multiplying the plasma levels of sildenafil, increasing the incidence and severity of side-effects. It is recommended that patients using protease inhibitors limit their use of sildenafil to no more than one 25-mg dose every 48 hours.

Some sildenafil users have complained of blurriness and loss of peripheral vision. In May of 2005, the U.S. Food and Drug Administration found that sildenafil could lead to vision impairment and a number of studies have linked sildenafil use with nonarteritic anterior ischemic optic neuropathy. When used with an alpha blocker, take them at least four hours apart to avoid hypotension.

[Home](#)|[Specials](#)|[New Products](#)|[All Products ...](#)|[Reviews](#)|[Contact Us](#)|[FAQ](#)|[Map](#)

BUY SILDENAPLEX 100 Copyright © 2005-2007. [Privacy Notice](#)